

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/130612>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

# Focus on post-resuscitation care

---

Claudio Sandroni (1), Alain Combes (2) and Jerry P. Nolan (3)

- (1) Istituto Anestesiologia e Rianimazione Università Cattolica del Sacro Cuore - Fondazione Policlinico Universitario "Agostino Gemelli" IRCCS. [claudio.sandroni@policlinicogemelli.it](mailto:claudio.sandroni@policlinicogemelli.it)
- (2) Sorbonne Université, INSERM, UMRS\_1166-ICAN, Institute of Cardiometabolism and Nutrition, and Service de médecine intensive-réanimation, Institut de Cardiologie, APHP Hôpital Pitié-Salpêtrière, F-75013 PARIS, France. [combes001@gmail.com](mailto:combes001@gmail.com)
- (3) Department of Anaesthesia, Royal United Hospital, Bath, and Warwick Medical School, University of Warwick, Coventry, CV4 7A, United Kingdom. [jerry.nolan@nhs.net](mailto:jerry.nolan@nhs.net)

**Key words:** cardiac arrest; cerebral perfusion pressure; coronary angiography; hypoxia-ischemia, brain; magnetic resonance imaging; mechanical ventilation; near-infrared spectroscopy; regional cerebral oxygenation; targeted temperature management.

**Word count:** 1050; **References:** 13; **Tables** 1

## ***Corresponding Author***

Claudio Sandroni  
Istituto Anestesiologia e Rianimazione Università Cattolica del Sacro Cuore  
Fondazione Policlinico Universitario "Agostino Gemelli" IRCCS  
Largo Francesco Vito, 1 – 00168 Rome, Italy  
email: [claudio.sandroni@policlinicogemelli.it](mailto:claudio.sandroni@policlinicogemelli.it)

In patients who are admitted to an intensive care unit after resuscitation from cardiac arrest (CA) mortality attains 60-70% [1]. Two-thirds of these deaths are due to hypoxic-ischaemic brain injury (HIBI) and occur mainly after 72 h, as a result of withdrawal of life sustaining treatment (WLST) based on prediction of a poor neurological outcome [2]. The second most common cause of post-resuscitation death, especially in the very early phase after return of spontaneous circulation (ROSC) is circulatory failure. Current research on post-resuscitation care focuses on interventions aimed at reducing these two important causes of morbidity and mortality after CA.

### Temperature management, coronary angiography

Targeted temperature management (TTM) for 24 hours after ROSC is recommended to reduce HIBI, but the optimal temperature is unclear [2, 3]. A recent multicentre pilot randomised clinical trial (RCT) assigned comatose survivors of witnessed out-of-hospital CA (OHCA) from ventricular fibrillation/ventricular tachycardia (VF/VT) to TTM at 32°C (n = 52), 33°C (n = 49) or 34°C (n = 49); see **Table 1**. Cooling was tolerated well, but 90-day neurological outcome did not differ among groups [4]. Patients cooled to 32°C had lower rates of WLST due to severe HIBI. Cooling to this temperature may be the focus of further research.

Immediate coronary angiography is recommended in survivors of OHCA with ST-segment elevation (STE) on their electrocardiogram [1], but the benefit of this intervention after non-STE OHCA is uncertain. The multicentre COACT trial, which included 552 non-STE patients resuscitated from VF/VT OHCA, showed that an immediate vs. a delayed angiography strategy (median [IQR] 2.3 [1.8–3.0] vs. 121.9 [52.0–197.3] hours from arrest) was not associated with higher 90-day survival (64.5 vs. 67.2%;  $p=0.51$ ) [5]. Acute unstable coronary lesions were found in less than 20% of the patients in this trial. Future studies may focus on non-STE patients more likely to benefit from immediate percutaneous coronary interventions (PCI). A recent retrospective study on 1,410 resuscitated OHCA also suggested that only patients with low risk of in-hospital death according to the Cardiac Arrest Hospital Prognosis score benefited from early PCI [6].

## Oxygenation and ventilation, cerebral perfusion

Observational evidence suggests that hyperoxia may be associated with worse neurological outcome after CA, while mild hypercapnia and higher mean arterial blood pressure (MAP) may be associated with better neurological outcome [2]. The COMACARE pilot multi-centre RCT evaluated the feasibility of targeting low-normal (4.5–4.7 kPa) vs. high-normal (5.8–6.0 kPa) PaCO<sub>2</sub>, normoxia (PaO<sub>2</sub> 10–15 kPa) vs. moderate hyperoxia (PaO<sub>2</sub> 20–25 kPa) and low-normal (65–75 mmHg) vs. high-normal (80–100 mmHg) MAP [7, 8] during the first 36 h after ROSC in 123 comatose patients resuscitated from VF/VT OHCA. The investigators achieved clear separation in PaO<sub>2</sub>, PaCO<sub>2</sub> and MAP between the groups but there was no difference in the values of neuron specific enolase (NSE), a biomarker of HIBI, at 48 h after ROSC. High-normal PaCO<sub>2</sub> and moderate hyperoxia increased regional cerebral oxygenation (rSO<sub>2</sub>) but there were no differences between the groups in any of the secondary outcomes.

One of several reasons for the failure to replicate the findings of previous observational studies is the likely considerable heterogeneity in optimal targets among patients. Approximately a third of post-CA patients will lose cerebral autoregulation and are likely to require a higher MAP to preserve cerebral blood flow [9]. rSO<sub>2</sub> has been used to determine optimal MAP in post-CA patients but it is unknown how well rSO<sub>2</sub> correlates with cerebral perfusion pressure (CPP). Taccone and colleagues have used trans-cranial Doppler to assess middle cerebral artery mean flow velocities (MFVmca) in 30 post-CA patients to provide an estimate of CPP (eCPP) [10]. Lower eCPP values were found in non-survivors compared with survivors, while MFVmca, rSO<sub>2</sub> and MAP values were similar between groups. It is possible that strategies to optimize eCPP instead of MAP or rSO<sub>2</sub> may improve neurological outcome.

## Prognostication

Prediction of neurological outcome in comatose resuscitated patients is challenging. Among indices for estimating severity of HIBI, NSE has several advantages: potential for blinded evaluation –

therefore avoiding a self-fulfilling prophecy bias – and no influence from sedation and TTM.

However, in order to obtain a specificity close to 100% for prediction of poor outcome, very high levels are needed, and the consequent sensitivity is low. Moreover, NSE values are influenced by haemolysis [1].

In a substudy of the TTM trial, [11] a new biomarker - neurofilament light chain (NFL) – was assessed blindly for neuroprognostication after CA. At 24 h, a NFL value of 478 pg/mL predicted poor neurological outcome at 6 months with 69 [57-79]% sensitivity and 98 [96-100]% specificity. The area under the receiver operating characteristic curve was 0.94 (0.92–0.95), greater than that of any other predictor, including NSE, brain CT, somatosensory evoked potentials, EEG, and clinical examination. Unlike NSE, NFL blood values are not influenced by haemolysis.

Clinical examination is crucial for neuroprognostication and absence of standard pupillary light reflex (sPLR) at  $\geq 72$  h is recommended as a robust predictor of poor neurological outcome after CA [1].

However, sPLR is operator-dependent, qualitative and, since it cannot be concealed from the treating team, is prone to self-fulfilling prophecy bias. In a European 456-patient multicentre prognostication study [12] where the quantitative neurological pupil index (NPi) measured blindly using automated pupillometry was compared with sPLR, an  $\text{NPi} \leq 2$  during the first 72h after ROSC had significantly higher specificity than an absent sPLR ( $p < 0.001$ ). NPi predicted unfavourable outcome with 0% FPR as early as 24 h after CA. Among patients whose sPLR was scored as absent, those with false positives had a significantly smaller pupil size than those who had a poor neurological outcome ( $1.9 \pm 0.22$  vs.  $2.8 \pm 1.05$  mm;  $p = 0.009$ ), suggesting that miosis may have prevented clinicians from detecting sPLR.

Neuroprognostication guidelines focus on the early post-resuscitation phase (72-96 h), while little is known about late predictors. In a multicentre prognostication study [13] fractional anisotropy (WWM-FA) measured on magnetic resonance imaging (MRI) was evaluated as a predictor of poor neurological outcome at 6 months in post-ROSC patients who had been unconscious for at least 7 days. A WWM-FA threshold of 0.91 was identified in a 150-patient derivation cohort as a prediction

and subsequently tested in a 50-patient validation cohort, where it showed 89.7 (75.8–97.1)% sensitivity and 100 (69.1–100)% specificity for prediction of poor outcome. Despite its complex measurement, WWM-FA is a promising predictor in late awakeners after CA.

## References

1. Nolan JP, Soar J, Cariou A, Cronberg T, Moulaert VR, Deakin CD, Bottiger BW, Friberg H, Sunde K, Sandroni C, European Resuscitation C, European Society of Intensive Care M (2015) European Resuscitation Council and European Society of Intensive Care Medicine 2015 guidelines for post-resuscitation care. *Intensive Care Med* 41: 2039-2056.
2. Nolan JP, Berg RA, Callaway CW, Morrison LJ, Nadkarni V, Perkins GD, Sandroni C, Skrifvars MB, Soar J, Sunde K, Cariou A (2018) The present and future of cardiac arrest care: international experts reach out to caregivers and healthcare authorities. *Intensive Care Med* 44: 823-832.
3. Aneman A, Cariou A, Nolan JP (2018) Understanding temperature goals after cardiac arrest. *Intensive Care Med* 44: 940-943.
4. Lopez-de-Sa E, Juarez M, Armada E, Sanchez-Salado JC, Sanchez PL, Loma-Orsorio P, Sionis A, Monedero MC, Martinez-Selles M, Martin-Benitez JC, Ariza A, Uribarri A, Garcia-Acuna JM, Villa P, Perez PJ, Storm C, Dee A, Lopez-Sendon JL (2018) A multicentre randomized pilot trial on the effectiveness of different levels of cooling in comatose survivors of out-of-hospital cardiac arrest: the FROST-I trial. *Intensive Care Med* 44: 1807-1815.
5. Lemkes JS, Janssens GN, van der Hoeven NW, Jewbali LSD, Dubois EA, Meuwissen M, Rijpstra TA, Bosker HA, Blans MJ, Bleeker GB, Baak R, Vlachojannis GJ, Eikemans BJW, van der Harst P, van der Horst ICC, Voskuil M, van der Heijden JJ, Beishuizen A, Stoel M, Camaro C, van der Hoeven H, Henriques JP, Vlaar APJ, Vink MA, van den Bogaard B, Heestermans T, de Ruijter W, Delnoij TSR, Crijns H, Jessurun GAJ, Oemrawsingh PV, Gosselink MTM, Plomp K, Magro M, Elbers PWG, van de Ven PM, Oudemans-van Straaten HM, van Royen N (2019) Coronary

- Angiography after Cardiac Arrest without ST-Segment Elevation. *N Engl J Med* 380: 1397-1407.
6. Bougouin W, Dumas F, Karam N, Maupain C, Marijon E, Lamhaut L, Jost D, Geri G, Beganton F, Varenne O, Spaulding C, Jouven X, Cariou A (2018) Should We Perform an Immediate Coronary Angiogram in All Patients After Cardiac Arrest?: Insights From a Large French Registry. *JACC Cardiovasc Interv* 11: 249-256.
  7. Jakkula P, Pettila V, Skrifvars MB, Hastbacka J, Loisa P, Tiainen M, Wilkman E, Toppila J, Koskue T, Bendel S, Birkelund T, Laru-Sompa R, Valkonen M, Reinikainen M, group Cs (2018) Targeting low-normal or high-normal mean arterial pressure after cardiac arrest and resuscitation: a randomised pilot trial. *Intensive Care Med* 44: 2091-2101.
  8. Jakkula P, Reinikainen M, Hastbacka J, Loisa P, Tiainen M, Pettila V, Toppila J, Lahde M, Backlund M, Okkonen M, Bendel S, Birkelund T, Pulkkinen A, Heinonen J, Tikka T, Skrifvars MB, group Cs (2018) Targeting two different levels of both arterial carbon dioxide and arterial oxygen after cardiac arrest and resuscitation: a randomised pilot trial. *Intensive Care Med* 44: 2112-2121.
  9. Sandroni C, Parnia S, Nolan JP (2019) Cerebral oximetry in cardiac arrest: a potential role but with limitations. *Intensive Care Med* *epub ahead of print*
  10. Taccone FS, Crippa IA, Creteur J, Rasulo F (2018) Estimated cerebral perfusion pressure among post-cardiac arrest survivors. *Intensive Care Med* 44: 966-967.
  11. Moseby-Knappe M, Mattsson N, Nielsen N, Zetterberg H, Blennow K, Dankiewicz J, Dragancea I, Friberg H, Lilja G, Insel PS, Rylander C, Westhall E, Kjaergaard J, Wise MP, Hassager C, Kuiper MA, Stammet P, Wanscher MCJ, Wetterslev J, Erlinge D, Horn J, Pellis T, Cronberg T (2019) Serum Neurofilament Light Chain for Prognosis of Outcome After Cardiac Arrest. *JAMA Neurol* 76: 64-71.
  12. Oddo M, Sandroni C, Citerio G, Miroz JP, Horn J, Rundgren M, Cariou A, Payen JF, Storm C, Stammet P, Taccone FS (2018) Quantitative versus standard pupillary light reflex for early

prognostication in comatose cardiac arrest patients: an international prospective multicenter double-blinded study. *Intensive Care Med* 44: 2102-2111.

13. Velly L, Perlberg V, Boulier T, Adam N, Delphine S, Luyt C-E, Battisti V, Torkomian G, Arbelot C, Chabanne R, Jean B, Di Perri C, Laureys S, Citerio G, Vargiolu A, Rohaut B, Bruder N, Girard N, Silva S, Cottenceau V, Tourdias T, Coulon O, Riou B, Naccache L, Gupta R, Benali H, Galanaud D, Puybasset L, Constantin J-M, Chastre J, Amour J, Vezinet C, Rouby J-J, Raux M, Langeron O, Degos V, Bolgert F, Weiss N, Similowski T, Demoule A, Duguet A, Tollard E, Veber B, Lotterie J-A, Sanchez-Pena P, Génestal M, Patassini M (2018) Use of brain diffusion tensor imaging for the prediction of long-term neurological outcomes in patients after cardiac arrest: a multicentre, international, prospective, observational, cohort study. *The Lancet Neurology* 17: 317-326.